

***Tetranychus urticae* literature**

Aguilar-Fenollosa, E., Pina, T., Gomez-Martinez, M.A., Hurtado, M.A. and Jacas, J.A. (2012) Does host adaptation of *Tetranychus urticae* populations in clementine orchards with a *Festuca arundinacea* cover contribute to a better natural regulation of this pest mite?. *Entomologia Experimentalis et Applicata*, 144: 181-190.

Aguilar-Fenollosa, E., Rey-Caballero, J., Blasco, J.M., Segarra-Moragues, J.G., Hurtado, M.A. and Jaques, J.A. (2016) Patterns of ambulatory dispersal in *Tetranychus urticae* can be associated with host plant specialization. *Experimental and Applied Acarology*, 68: 1-20.

Albendin, G., Garcia, M.D. and Molina, J.M. (2015) Multiple natural enemies do not improve two spotted spider mite and flower western thrips control in strawberry tunnels. *Chilean Journal of Agricultural Research*, 75: 63-70.

Alekseev, Y.I., Daricheva, M.A., Zavodchikova, V.V., Kamalov, K., Niyazov, O.D. and Sukhoruchenko, G.I. (1976) The arthropod fauna of cotton in the Murgab oasis. Ecology and economic importance of the insects of Turkmenia.: *Ekologiya i khozyaistvennoe znachenie nasekomykh Turkmenii*. Ylim, Ashkhabad, USSR,.

Altincag, R. and Akten, T. (1993) Insect pests in grapevine nurseries and remedies in Aegean region: problems and their solutions. *Bitki Koruma Bulteni*, 33: 153-165.

Alvarez, J.M., Bellotti, A.C., Braun, A.R. and Acosta, A. (1991) Studies of the pathogenicity of a fungus associated with *Tetranychus urticae* Koch, a pest of cassava. *Revista Colombiana de Entomología*, 17: 28-33.

Alvarez, L.D. and Alarcon, L. (2004) Catalogo de caros fitofagos de la provincia de Las Tunas. *Fitosanidad*, 8: 19-21.

Amini, M.Y., Ullah, M.S., Kitagawa, A., Kanazawa, R., Takano, Y., Suzuki, T. and Gotoh, T. (2016) Scotophase interruption with LEDs and OLEDs to inhibit photoperiodic induction of diapause in *Tetranychus urticae* and *T. kanzawai* (Acari: Tetranychidae). *Systematic and Applied Acarology*, 21: 1436.

André, M. (1945) Sur les dommages causés en France aux culture de soja par l'invasion d'un Tétranyque. *Comptes rendus de l'Académie d'Agriculture de France*, 1-2.

Andrews, K.L. and Barnes, M.M. (1981) Spider mites on almond in the southern San Joaquin Valley of California. *Environmental Entomology*, 10: 6-9.

Andrews, K.L. and Poe, S.L. (1980) Spider mites of El Salvador, Central America (Acari: Tetranychidae). *Florida Entomologist*, 63: 502-505.

Aranda, B.R. (1974) Tetranychoidea (Acari) de uma area de cerrado do estado de São Paulo. Escola Superior de Agricultura "Luiz de Queiroz", Piracicaba, São Paulo: 47.

Arias Giralda, A. and Nieto Calderon, J. (1983) Estimacion de las perdidas producidas por la "araña amarilla" (*Tetranychus urticae* Koch) en Tierra de Barros (Badajoz) y propuesta de un umbral de tolerancia economica. Boletín del Servicio de Plagas, 9: 227-252.

Arnemann, J.A., Fiorin, R.A., Perini, C.R., Storck, L., Curioletti, L.E., Nachman, G. and Guedes, J.V.C. (2015) Density and growth rates of spider mites in relation to phenological stages of soybean cultivars in Brazil. Experimental and Applied Acarology, 67: 423-440.

Arthur, V., Nicastro, R.L., Sato, M.E. and Machi, A.R. (2016) Milbemectin and etoxazol acaricide resistant and susceptible strains of *Tetranychus urticae* (Trombidiformes: Tetranychidae) are equally radiosusceptible and unable to reproduce when irradiated with 400 Gy. Florida Entomologist, 99: 34-37.

Artofer, R. (1976) Spider-mites in vine cultivation. Pflanzenarzt, 29: 73.

Atanasov, N. (1974) Experiments on biological control of *Tetranychus urticae* Koch and *Tetranychus turkestani* Ugarov et Nikolski in Bulgaria. Proceedings of the 4th International Congress of Acarology, Akademiai Kiado, Budapest, Hungary: 663-666.

Baillod, M., Antonin, P. and Mittaz, C. (1989) Migrations, estimation des populations et nuisibilité de l'acarien jaune commun, *Tetranychus urticae* Koch dans la viticulture valaisanne. Revue Suisse de Viticulture, d'Arboriculture et d'Horticulture, 21: 179-183.

Baker, E.W. (1975) Plant-feeding mites of Thailand (Tetranychidae, Tenuipalpidae, and Tuckerellidae). Plant Protection Service Technical Bulletin. Bangkok, Thailand.,

Baker, E.W. and Pritchard, A.E. (1960) The tetranychoid mites of Africa. Hilgardia, 29: 455-574.

Barbar, Z. (2016) The mite fauna (Acari) of two Syrian citrus orchards, with notes on their morphology and economic importance. Systematic & Applied Acarology, 21: 991-1008.

Basu, B.D. (1963) Studies on two new mites of the family Tetranychidae (genus *Tetranychus*). Proceedings of the Zoological Society of Calcutta, 16: 221-224.

Bayu, M., Ullah, M.S., Takano, Y. and Gotoh, T. (2017) Impact of constant versus fluctuating temperatures on the development and life history parameters of *Tetranychus urticae* (Acari: Tetranychidae). Experimental and Applied Acarology, 72: 205-227.

Beer, R.E. and Lang, D.S. (1958) The Tetranychidae of Mexico. University of Kansas Scientific Bulletin, 38: 1231-1259.

Bellini, M.R., Moraes, G.J.d. and Feres, R.J.F. (2005) Acaros (Acari) de dois sistemas de cultivo de seringueira no Noroeste do Estado de São Paulo. Neotropical Entomology, 34: 475-484.

Ben-David, T., Melamed, S., Gerson, U. and Morin, S. (2007) ITS2 sequences as barcodes for identifying and analyzing spider mites (Acari: Tetranychidae). Experimental & Applied Acarology, 41: 169-181.

Bi, J.L., Niu, Z.M., Yu, L. and Toscano, N.C. (2016) Resistance status of the carmine spider mite, *Tetranychus cinnabarinus* and the twospotted spider mite, *Tetranychus urticae* to selected acaricides on strawberries. *Insect Science*, 23: 88-93.

Bibby, F.F. and Tuttle, D.M. (1959) Notes on phytophagous and predatory mites of Arizona. *Journal of Economic Entomology*, 52: 186-190.

Bolland, H.R. (1999) Verslag Baikalreis 16 juli t:m 3 augustus 1999. Vereniging voor veldbiologie, 1999,.

Breeuwer, J.A.J. (1997) Wolbachia and cytoplasmic incompatibility in the spider mites *Tetranychus urticae* and *T. turkestanii*. *Heredity*, 79: 41-47.

Bugeme, D.M., Knapp, M., Boga, H.I., Ekesi, S. and Maniania, N.K. (2014) Susceptibility of developmental stages of *Tetranychus urticae* (Acari: Tetranychidae) to infection by *Beauveria bassiana* and *Metarhizium anisopliae* (Hypocreales: Clavicipitaceae). *International Journal of Tropical Insect Science*, 34: 190-196.

Carbonnelle, S., Hance, T., Migeon, A., Baret, P., Cros-Arteil, S. and Navajas, M. (2007) Microsatellite markers reveal spatial genetic structure of *Tetranychus urticae* (Acari: Tetranychidae) populations along a latitudinal gradient in Europe. *Experimental & Applied Acarology*, 41: 225-241.

Carmona, M.M. (1960) Contribuição para o conhecimento dos ácaros das plantas cultivadas em Portugal - I. *Agronomia Lusitana*, 22: 221-230.

Carmona, M.M. (1967) Contribuição para o estudo de alguns ácaros fitófagos e depredadores, de Angola. *Agronomia Lusitana*, 29: 267-288.

Carmona, M.M. (1970) Contribuição para o conhecimento dos ácaros das plantas cultivadas em Portugal - V. *Agronomia Lusitana*, 31: 137-183.

Carmona, M.M. (1973) Acaros fitofagos e predadores da Ilha da Madeira. *Agronomia Lusitana*, 34: 255-281.

Chaudhri, W.M., Akbar, S. and Rasool, A. (1974) Taxonomic studies of the mites belonging to the families Tenuipalpidae, Tetranychidae, Tuckerellidae, Caligonellidae, Stigmeidae and Phytoseiidae - PL-480 Project on mites. Lyallpur, Pakistan, University of Agriculture: 250 p.

Chen, D.-S., Jin, P.-Y., Zhang, K.-J., Ding, X.-L., Yang, S.-X., Ju, J.-F., Zhao, J.-Y. and Hong, X.-Y. (2014) The complete mitochondrial genomes of six species of *Tetranychus* provide insights into the phylogeny and evolution of spider mites. *Plos One*, 9: e110625.

Cheng, G., Ding, X.L., Zhang, J.P. and Hong, X.Y. (2013) *Tetranychus urticae* (green form) on *Gossypium hirsutum* in China: two records confirmed by aedeagus morphology and RFLP analysis. *Systematic and Applied Acarology*, 18: 239-244.

Choi, O., Park, J.J. and Kim, J. (2016) *Tetranychus urticae* (Acari: Tetranychidae) transmits Acidovorax citrulli, causal agent of bacterial fruit blotch of watermelon. *Experimental and Applied Acarology*, 69: 445-451.

Clemente, S.H., Rodrigues, L.R., Ponce, R., Varela, S.A.M. and Magalhaes, S. (2016) Incomplete species recognition entails few costs in spider mites, despite first-male precedence. *Behavioral Ecology and Sociobiology*, 70: 1161-1170.

Corpuz-Raros, L.A. (2001) New mite pests and new host records of phytophagous mites (Acari) from the Philippines. *Philippine Agricultural Scientist*, 84: 341-351.

Costa Comelles, J., Soares, A.O., Vercher, R., Schanderl, H. and Ferragut, F. (1993) A contribution to data on mite fauna in the Azores Islands. *elatórios e Comunicações do Departamento de Biologia da Universidade dos Açores*, 22: 40-44.

Cruz, W.P.d., Sarmento, R.A., Teodoro, A.V., Erasmo, E.A.L., Pedro Neto, M., Ignacio, M. and Ferreira Junior, D.F. (2012) Acarofauna em cultivo de pinhão-manso e plantas espontâneas associadas. *Pesquisa Agropecuaria Brasileira*, 47: 319-327.

Cuenod, H. (1956) Les acariens nuisibles aux agrumes de Tunisie. 4th International Congress of Mediterranean Agrumiculture, Tel-Avis, Israel : may 20-27th 1956: 299-305.

de Castro, T.R., Wekesa, V.W., Moral, R.D., Demetrio, C.G.B., Delalibera, I. and Klingen, I. (2013) The effects of photoperiod and light intensity on the sporulation of Brazilian and Norwegian isolates of *Neozygites floridana*. *Journal of Invertebrate Pathology*, 114: 230-233.

de Villiers, M. and Pringle, K.L. (2011) The presence of *Tetranychus urticae* (Acari: Tetranychidae) and its predators on plants in the ground cover in commercially treated vineyards. *Experimental and Applied Acarology*, 53: 121-137.

Dobosz, R., Skorupska, A. and Blaszak, C. (1995) The appearance of spider mites (Tetranychidae) in parks of Poznan. Boczek, J. and Ignatowicz, S., Materiały z Sympozjum na temat: "Osiagniecia Akarologii w Polsce", Siedlce, 26-27 września 1995, Komitet Ochrony Roslin Polska Akademia Nauk: 39-42.

Domingos, C.A., Melo, J.W.S., Oliveira, J.E.M. and Gondim, M.G.C. (2014) Mites on grapevines in northeast Brazil: occurrence, population dynamics and within-plant distribution. *International Journal of Acarology*, 40: 145-151.

Dube, Z.P., Mashela, P.W. and Abdelgadir, A.H. (2015) Identification of spider mites from *Moringa oleifera* using molecular techniques. *Acta Agriculturae Scandinavica Section B-Soil and Plant Science*, 65: 479-482.

Ehara, S. (1956) Notes on some tetranychid mites of Japan (with 37 txt-figures). *Japanese Journal of Applied Zoology*, 21: 139-147.

Ehara, S. (1966) Some mites associated with plants in the State of São Paulo, Brazil, with a list of plant mites of South America. *Japanese Journal of Zoology*, 15: 129-150.

Elbadry, E.A. and Kansouh, A.S. (1975) On the population dynamics of the spider mite *Tetranychus cinnabarinus* (Boisd.) and the predatory mite *Amblyseius gossypii* (Elbadry) on cotton. *Anzeiger für Schadlingskunde Pflanzenschutz Umweltschutz*, 48: 35-37.

Emmanouel, N.G. and Papadoulis, G.T. (1987) *Panonychus citri* (MacGregor) (Tetranychidae) and *Eriophyes medicaginis* K. (Eriophyidae): two important phytophagous mites recorded for the first time in Greece. *Entomologia Hellenica*, 5: 3-6.

Estebanes-Gonzalez, M.L. and Baker, E.W. (1968) Arañas rojas de Mexico (Acarina: Tetranychidae).. Anales de la Escuela Nacional de Ciencias Biologicas, 15: 61-133.

Estebanes-Gonzalez, M.L. and Rodriguez-Navarro, S. (1991) Observaciones sobre algunos acaros de las familias Tetranychidae, Eriophyidae, Acaridae y Tarsonemidae (Acari), en hortalizas de Mexico. *Folia Entomologica Mexicana*, 83: 199-212.

Ferragut, F. and Santonja, M.C. (1989) Taxonomia y distribucion de los acaros del genero *Tetranychus* Dufour 1832 (Acari: Tetranychidae), en Espana. *Boletin de Sanidad Vegetal, Plagas*, 15: 271-281.

Ferreira, C.B.S., Andrade, F.H.N., Rodrigues, A.R.S., Siqueira, H.A.A. and Gondim, M.G.C. (2015) Resistance in field populations of *Tetranychus urticae* to acaricides and characterization of the inheritance of abamectin resistance. *Crop Protection*, 67: 77-83.

Ferrero, M., Moraes, G.J.d., Kreiter, S., Tixier, M.S. and Knapp, M. (2007) Life tables of the predatory mite *Phytoseiulus longipes* feeding on *Tetranychus evansi* at four temperatures (Acari: Phytoseiidae, Tetranychidae). *Experimental & Applied Acarology*, 41: 45-53.

Fiaboe, K.K.M., Gondim, M.G.C.J., Moraes, G.J.d., Ogol, C.K.P.O. and Knapp, M. (2007) Surveys for natural enemies of the tomato red spider mite *Tetranychus evansi* (Acari: Tetranychidae) in northeastern and southeastern Brazil. *Zootaxa*, 1395: 33-58.

Flechtmann, C.H.W. (1996) Rediscovery of *Tetranychus abacae* Baker & Pritchard, additional description and notes on South American spider mites (Acari, Prostigmata, Tetranychidae). *Revista Brasileira de Zoologia*, 13: 569-578.

Flechtmann, C.H.W. and Abreu, J.M. (1973) Acaros fitofagos do Estado da Bahia, Brasil. (Notas preliminares). *Ciencia e Cultura*, 25: 244-251.

Flechtmann, C.H.W. and Bastos, J.A.M. (1972) Acaros Tetranychoidea do estado do Ceara, Brasil. *Ciencia e Agrotecnologia*, 2: 83-90.

Flechtmann, C.H.W. and Etienne, J. (2006) Further notes on plant associated mites (Acari) from Guadeloupe and Les Saintes, lesser Antilles. *International Journal of Acarology*, 32: 377-382.

Flechtmann, C.H.W. and Hunter, P.E. (1971) The spider mites (Prostigmata : Tetranychidae) of Georgia. *Journal of the Georgia Entomological Society*, 6: 16-30.

Flechtmann, C.H.W., Kreiter, S., Etienne, J. and Moraes, G.J.d. (1999) Plant mites (Acari) of the French Antilles. 1. Tetranychoidea (Prostigmata). *Acarologia*, 40: 137-144.

Flores, J.L., Chavez, E.C., Uribe, L.A.A., Canales, R.F. and Fuentes, Y.M.O. (2013) Demographic parameters of *Tetranychus urticae* (Acari: Tetranychidae) on four Rosa sp cultivars. *Florida Entomologist*, 96: 1508-1512.

Froggatt, W.W. (1921) Orchard and garden mites. N° 2 - Spinning mites (Family Tetranychidae). Agricultural Gazette of New South Wales, 32: 130-130.

Funayama, K. (2015) Outbreaks of the two-spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae) are caused by broad-spectrum insecticide spraying in apple orchards. Applied Entomology and Zoology, 50: 169-174.

Funayama, K., Komatus, M., Sonoda, S., Takahashi, I. and Hara, K. (2015) Management of apple orchards to conserve generalist phytoseiid mites suppresses two-spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae). Experimental and Applied Acarology, 65: 43-54.

Furtado, I.P., Moraes, G.J.d., Kreiter, S. and Knapp, M. (2006) Search for effective natural ennemis of *Tetranychus evansi* in south and southeast Brazil. Experimental & Applied Acarology, 40: 157-174.

Furtado, I.P., Toledo, S., Moraes, G.J.d., Kreiter, S. and Knapp, M. (2007) Search for effective natural enemies of *Tetranychus evansi* (Acari: Tetranychidae) in northwest Argentina. Experimental and Applied Acarology, 43: 121-127.

Fytrou, N. and Tsagkarakou, A. (2014) Reproductive incompatibility between genetically differentiated populations of *Tetranychus urticae* from different host plants. Entomologia Hellenica, 23: 1-10.

Ghasemi Moghadam, S., Ahadiyat, A. and Ueckermann, E.A. (2016) Species composition of tetranychoid mites (Acari: Trombidiformes: Prostigmata: Tetranychidae) in main landscapes of Tehran and modelling ecological niche of Tetranychidae in main climates of Tehran Province, Iran. Biologia, 71: 1151-1166.

Ghazy, N.A., Otsuki, H., Sekido, T., Yano, S. and Amano, H. (2016) Dispersal of diapausing *Tetranychus urticae* and *T. kanzawai*. Entomologia Experimentalis Et Applicata, 160: 126-132.

Giertych, M.J., Karolewski, P. and Oleksyn, J. (2008) Early-season defoliation of *Sorbus aucuparia* (L.) and *Acer platanoides* (L.) can induce defense mechanisms againts the spider mite *Tetranychus urticae* (Koch). Polish Journal of Ecology, 56: 443-452.

Goka, K. and Takafuji, A. (1995) Allozyme variations among populations of the two-spotted spider mite, *Tetranychus urticae* Koch, in Japan. Applied Entomology & Zoology, 30: 567-579.

Golizadeh, A., Ghavidel, S., Razmjou, J., Fathi, S.A.A. and Hassanpour, M. (2017) Comparative life table analysis of *Tetranychus urticae* Koch (Acari: Tetranychidae) on ten rose cultivars. Acarologia, 57: 607-616.

González-Zamora, J., López, C. and Avilla, C. (2011) Population studies of arthropods on *Melia azedarach* in Seville (Spain), with special reference to *Eutetranychus orientalis* (Acari: Tetranychidae) and its natural enemies. Experimental and Applied Acarology, 55: 389-400.

Gotoh, T. (1997) Annual life cycles of populations of the two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae) in four Japanese pear orchards. Applied Entomology & Zoology, 32: 207-216.

Gotoh, T. and Tokioka, T. (1996) Genetic compatibility among diapausing red, non-diapausing red and diapausing green forms of the two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae). Japanese Journal of Entomology, 64: 215-225.

Gotoh, T., Bruin, J., Sabelis, M.W. and Menken, S.B.J. (1993) Host race formation in *Tetranychus urticae*: genetic differentiation, host plant preference, and mate choice in a tomato and a cucumber strain. Entomologia Experimentalis et Applicata, 68: 171-178.

Gotoh, T., Kitashima, Y. and Goka, K. (2007) *Tetranychus* mite species identification using esterase and phosphoglucomutase zymograms. Applied Entomology and Zoology, 42: 579-585.

Gotoh, T., Kitashima, Y. and Sato, T. (2013) Effect of hot-water treatment on the two-spotted spider mite, *Tetranychus urticae*, and its predator, *Neoseiulus californicus* (Acari: Tetranychidae, Phytoseiidae). International Journal of Acarology, 39: 533-537.

Gotoh, T., Moriya, D. and Nachman, G. (2015) Development and reproduction of five *Tetranychus* species (Acari: Tetranychidae): Do they all have the potential to become major pests?. Experimental and Applied Acarology, 66: 453-479.

Gotoh, T., Noda, H. and Hong, X.Y. (2003) Wolbachia distribution and cytoplasmic incompatibility based on a survey of 42 spider mite species (Acari: Tetranychidae) in Japan. Heredity, 91: 208-216.

Gotoh, T., Noda, H. and Ito, S. (2007) Cardinium symbionts cause cytoplasmic incompatibility in spider mites. Heredity, 98: 13-20.

Gotoh, T., Saito, M., Suzuki, A. and Nachman, G. (2014) Effects of constant and variable temperatures on development and reproduction of the two-spotted spider mite *Tetranychus urticae* (Acari: Tetranychidae). Experimental and Applied Acarology, 64: 465-478.

Gotoh, T., Sugasawa, J. and Nagata, T. (1999) Reproductive compatibility of the two-spotted spider mite (*Tetranychus urticae*) infected with Wolbachia. Entomological Science, 2: 289-295.

Gotoh, T., Sugasawa, J., Noda, H. and Kitashima, Y. (2007) Wolbachia-induced cytoplasmic incompatibility in Japanese populations of *Tetranychus urticae* (Acari: Tetranychidae). Experimental & Applied Acarology, 41: 1-16.

Gotoh, T., Takafuji, A. and Gomi, K. (1996) Tetranychid mites of Okinawa Island (Acari: Tetranychidae). Journal of the Acarological Society of Japan, 5: 89-97.

Guanilo, A.D., de Moraes, G.J., Flechtmann, C.H.W. and Knapp, M. (2012) Phytophagous and fungivorous mites (Acari: Prostigmata, Astigmata) from Peru. International Journal of Acarology, 38: 120-134.

Gugole Ottaviano, M.F., Sanchez, N.E., Roggiero, M.F. and Greco, N.M. (2013) Performance of *Tetranychus urticae* and *Neoseiulus californicus* on strawberry cultivars and assessment of the effect of glandular trichomes. Arthropod-Plant Interactions, 7: 547-554.

Gupta, S.K. (1992) Arachnida: plant mites (Acari). Zoological Survey of India, State Fauna Series 3 : Fauna of West Bengal, Part 3: 61-211.

Gupta, S.K. (1995) Plant mites (Acari). Zoological Survey of India, State Fauna Series 4 : Fauna of Meghalaya, Part 2: 17-50.

Gutierrez, J. (1974) Les espèces du genre *Tetranychus* Dufour (Acariens : Tetranychidae) ayant une incidence économique à Madagascar et dans les îles voisines. Compétition entre les complexes *Tetranychus neocaledonicus* André et *Tetranychus urticae* Koch. *Acarologia*, 16: 258-270.

Gutierrez, J. and Etienne, J. (1981) Quelques données sur les acariens Tetranychidae attaquant les plantes cultivées au Sénégal. *Agronomie Tropicale*, 36: 391-394.

Gutierrez, J. and Etienne, J. (1986) Les Tetranychidae de l'île de la Réunion et quelques-uns de leurs prédateurs. *Agronomie Tropicale*, 41: 84-91.

Hada, H., Hinomoto, N. and Gotoh, T. (2016) Genetic structure of *Tetranychus urticae* (Acari: Tetranychidae) populations under acaricide selection pressure assessed using microsatellite markers. *Systematic and Applied Acarology*, 21: 878-888.

Harder, M.J., Tello, V.E. and Giliomee, J.H. (2016) The acaricidal effect of ethanolic extracts of *Chenopodium quinoa* Willd. on *Tetranychus urticae* Koch (Acari: Tetranychidae). *African Entomology*, 24: 50-60.

Harris, A.L., Ullah, R. and Fountain, M.T. (2017) The evaluation of extraction techniques for *Tetranychus urticae* (Acari: Tetranychidae) from apple (*Malus domestica*) and cherry (*Prunus avium*) leaves. *Experimental and Applied Acarology*, 72: 367-377.

Harvey, F.L. (1892) The two-spotted mite. Annual Report Maine Agricultural Experimental Station, 133-144.

Hata, F.T., Ventura, M.U., Carvalho, M.G., Miguel, A.L.A., Souza, M.S.J., Paula, M.T. and Zawadneak, M.A.C. (2016) Intercropping garlic plants reduces *Tetranychus urticae* in strawberry crop. *Experimental and Applied Acarology*, 69: 311-321.

Helle, W. (1965) Inbreeding depression in an arrhenotokous mite (*Tetranychus urticae* Koch). *Entomologia Experimentalis et Applicata*, 8: 299-304.

Hill, R.L. and O'Donnell, D.J. (1991) Reproductive isolation between *Tetranychus lintearius* and two related mites, *T. urticae* and *T. turkestanii* (Acarina: Tetranychidae). *Experimental & Applied Acarology*, 11: 241-251.

Hinomoto, N. and Takafuji, A. (2001) Genetic diversity and phylogeny of the Kanzawa spider mite, *Tetranychus kanzawai*, in Japan. *Experimental & Applied Acarology*, 25: 355-370.

Hinomoto, N. and Takafuji, A. (2004) Evaluation of mitochondrial cytochrome oxydase subunit I sequences in *Tetranychus kanzawai* Kishida (Acari: Tetranychidae) for phylogeographic studies. *Journal of the Acarological Society of Japan*, 13: 47-55.

Hinomoto, N., Dinh Pha, T., Anh Tuan, P., Thi Bao Ngoc, L., Tajima, R., Ohashi, K., Osakabe, M. and Takafuji, A. (2007) Identification of spider mites (Acari: Tetranychidae) by DNA sequences: a case study in northern Vietnam. *International Journal of Acarology*, 33: 53-60.

Hinomoto, N., Osakabe, M., Gotoh, T. and Takafuji, A. (2001) Phylogenetic analysis of green and red forms of the two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae), in Japan, based on mitochondrial cytochrome oxidase subunit I sequences. *Applied Entomology & Zoology*, 36: 459-464.

Hurtado, M.A., Ansaloni, T., Cros-Arteil, S., Jacas, J.A. and Navajas, M. (2008) Sequence analysis of the ribosomal internal transcribed spacers region in spider mites (Prostigmata: Tetranychidae) occurring in citrus orchards in Eastern Spain: use for species discrimination. *Annals of Applied Biology*, 153: 167-174.

Iacob, N. and Lefter, G. (1960) Observations sur l'expansion de quelques espèces d'acariens nuisibles dans les vergers de Roumanie. *Acarologia*, 2: 66-74.

Imura, O. (2003) Herbivorous arthropod community of an alien weed *Solanum carolinense* L. *Applied Entomology and Zoology*, 38: 293-300.

Iqbal, I. and Ali, A. (2008) Red spider mites from fruit orchards of Abbottabad, including a new species. *Biologia (Pakistan)*, 54: 125-130.

Iwassaki, L., Sato, M., Calegario, F., Poletti, M. and Maia, A. (2015) Comparison of conventional and integrated programs for control of *Tetranychus urticae* (Acari: Tetranychidae). *Experimental and Applied Acarology*, 65: 205-217.

Izadi, H., Asadabadi, A., Khanjani, M. and Payandeh, A. (2010) Some phytophagous mites associated with pomegranate, palm and citrus in southeast Iran. 1,, p. 112.. Moraes de, G.J., Castilho, R.C. and Flechtmann, C.H.W., 13th International Congress of Acarology, Recife-PE,Brazil, Abstract book: 112.

Jiao, R., Xu, C.X., Yu, L.C., He, X.Z., Qiao, G.Y., He, L.M. and Li, L.T. (2016) Prolonged coldness on eggs reduces immature survival and reproductive fitness in *Tetranychus urticae* (Acari: Tetranychidae). *Systematic and Applied Acarology*, 21: 1651-1661.

Karlec, F., Duarte, A.D.F., Oliveira, A.N.A.C.B.D.E. and Cunha, U.S.D. (2017) Development of *Tetranychus urticae* Koch (Acari: Tetranychidae) in different strawberry cultivars. *Revista Brasileira de Fruticultura*, 39: e-171.

Kasap, I., Polat, B. and Kok, S. (2014) The important pest and predatory mites species (Acari) and their population fluctuation in the vineyards of Canakkale Province. *Turkiye Entomoloji Dergisi-Turkish Journal of Entomology*, 38: 451-458.

Kawashima, M., Chung, B.K. and Jung, C.L. (2008) Herbivorous and predacious mites on persimmon trees, *Diospyros kaki* Thunb., in Korea. *International Journal of Acarology*, 34: 167-174.

Khalil-Manesh, B. (1973) Phytophagous mite fauna of Iran. *Entomologie et Phytopathologie Appliquées*, 30-38.

Kontschán, J. and Ripka, G. (2017) Checklist of the Hungarian spider mites and flat mites (Acari: Tetranychidae and Tenuipalpidae). *Systematic and Applied Acarology*, 22: 1199-1225.

- Kreiter, S., Auger, P., Lebdi Grissa, K., Tixier, M.S., Chermiti, B. and Dali, M. (2002) Plant inhabiting mites (Acari: Prostigmata & Mesostigmata) of some Northern Tunisian crops. *Acarologia*, 42: 389-402.
- Lamb, K.P. (1953) Survey of red spider mites (Acarina: Tetranychidae) on grape vines. *New Zealand Journal of Science and Technology (A)*, 35: 65-66.
- Lee, M., Suh, S. and Kwon, Y. (1999) Phylogeny and diagnostic markers of six *Tetranychus* species (Acarina: Tetranychidae) in Korea based on the mitochondrial cytochrome oxidase subunit I. *Journal of Asia Pacific Entomology*, 2: 85-92.
- Leite, G.L.D., Picanco, M., Zanuncio, J.C. and Marquini, F. (2003) Factors affecting mite herbivory on eggplants in Brazil. *Experimental and Applied Acarology*, 31: 243-252.
- Li, J. and Margolies, D.C. (1994) Barometric pressure influences initiation of aerial dispersal in the twospotted spider mite. *Journal of the Kansas Entomological Society*, 67: 386-393.
- Livshits, I.Z. and Salinas-Croche, A. (1968) Preliminares acerca de los acaros "Tetranicos" de Cuba. Cuba, Cent. Natl. Fitosenit.: 156 p.
- Lopes Ribeiro, A.E., Correa Gondim, M.G., Jr., Calderan, E. and Delalibera, I., Jr. (2009) Host range of *Neozygites floridana* isolates (Zygomycetes: Entomophthorales) to spider mites. *Journal of Invertebrate Pathology*, 102: 196-202.
- Lu, W., Wang, M., Xu, Z., Shen, G., Wei, P., Li, M., Reid, W. and He, L. (2017) Adaptation of acaricide stress facilitates *Tetranychus urticae* expanding against *Tetranychus cinnabarinus* in China. *Ecology and Evolution*, 7: 1233-1249.
- Lucini, T., Resende, J.T.V., Oliveira, J.R.F., Scabeni, C.J., Zeist, A.R. and Resende, N.C.V. (2016) Repellent effects of various cherry tomato accessions on the two-spotted spider mite *Tetranychus urticae* Koch (Acari: Tetranychidae). *Genetics and Molecular Research*, 15: 6.
- Maleknia, B., Fathipour, Y. and Soufbaf, M. (2016) How greenhouse cucumber cultivars affect population growth and two-sex life table parameters of *Tetranychus urticae* (Acari: Tetranychidae). *International Journal of Acarology*, 42: 70-78.
- Manson, D.C.M. (1963) Mites of the families Tetranychidae and Tenuipalpidae associated with citrus in South East Asia. *Acarologia*, 5: 351-364.
- Marcic, D. (2007) Sublethal effects of spirodiclofen on life history and life-table parameters of two-spotted spider mite (*Tetranychus urticae*). *Experimental & Applied Acarology*, 42: 121-129.
- Marcic, D., Petronijevic, S., Drobnjakovic, T., Prijovic, M., Peric, P. and Milenovic, S. (2012) The effects of spirotetramat on life history traits and population growth of *Tetranychus urticae* (Acari: Tetranychidae). *Experimental and Applied Acarology*, 56: 113-122.
- Marinosci, C., Magalhaes, S., Macke, E., Navajas, M., Carbonell, D., Devaux, C. and Olivier, I. (2015) Effects of host plant on life-history traits in the polyphagous spider mite *Tetranychus urticae*. *Ecol Evol*, 5: 3151-8.

Markoyiannaki-Printzioui, D., Papaioannou-Souliotis, P., Zeginis, G. and Giatropoulos, C. (2000) Observations on acarofauna in four apple orchards of Central Greece. I. Incidence of pedoclimatic conditions and agricultural techniques on phytoseiid mites (Acari: Phytoseiidae). *Acarologia*, 41: 109-126.

Martinez, A.Z., Torre de la , P.E. and Garcia, S.A. (2004) Principales acaros detectados en la provincia de Villa Clara. *Fitosanidad*, 8: 3-17.

Matsuda, T., Morishita, M., Hinomoto, N. and Gotoh, T. (2014) Phylogenetic analysis of the spider mite sub-family Tetranychinae (Acari: Tetranychidae) based on the mitochondrial COI gene and the 18S and the 5' end of the 28S rRNA genes indicates that several genera are polyphyletic. *Plos One*, 9: e108672.

Memarizadeh, N., Ghadamyari, M., Zamani, P. and Sajedi, R.H. (2013) Resistance mechanisms to abamectin in Iranian populations of the two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae). *Acarologia*, 53: 235-246.

Mendonca, R.S., Navia, D., Diniz, I.R., Auger, P. and Navajas, M. (2011) A critical review on some closely related species of *Tetranychus* sensu stricto (Acari: Tetranychidae) in the public DNA sequences databases. *Experimental and Applied Acarology*, 55: 1-23.

Mercado, V.T. and Arriagada, M.D. (2014) Preliminary studies of lethal and sub-lethal effects of ethanolic extracts of four xerophytic species of high Andes of Chilean against *Tetranychus cinnabarinus* (Acarina: Tetranychidae). *Revista De La Facultad De Ciencias Agrarias*, 46: 135-148.

Meyer, M.K.P.S. (1974) A revision of the Tetranychidae of Africa (Acari) with a key to the genera of the world. *Entomology Memoir*, Department of Agricultural Technical Services, Republic of South Africa, 1-291.

Mladenović, K., Stojnić, B., Vidović, B. and Radulović, Z. (2013) New records of the tribe Bryobiini berlsese (Acari: Tetranychidae: Bryobiinae) from Serbia, with notes about associated predators (Acari: Phytoseiidae). *Archives of Biological Sciences*, 65: 1199-1210.

Moghadam, M.M., Ghadamyari, M. and Talebi, K. (2012) Resistance mechanisms to fenazaquin in Iranian populations of two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae). *International Journal of Acarology*, 38: 138-145.

Monteiro, V.B., Gondim, M.G.C., Oliveira, J.E.D., Siqueira, H.A.A. and Sousa, J.M. (2015) Monitoring *Tetranychus urticae* Koch (Acari: Tetranychidae) resistance to abamectin in vineyards in the Lower Middle Sao Francisco Valley. *Crop Protection*, 69: 90-96.

Motahari, M., Kheradmand, K., Roustaei, A.M. and Talebi, A.A. (2014) The impact of cucumber nitrogen nutrition on life history traits of *Tetranychus urticae* (Koch) (Acari: Tetranychidae). *Acarologia*, 54: 443-452.

Najafabadi, S.S.M., Shoushtari, R.V., Zamani, A.A., Arbabi, M. and Farazmand, H. (2014) Life Parameters of *Tetranychus urticae* (Acari: Tetranychidae) on Six Common Bean Cultivars. *Journal of Economic Entomology*, 107: 614-622.

Nassar, O.A. and Ghai, S. (1981) Taxonomic studies on tetranychoid mites infesting vegetable and fruit crops in Delhi and surrounding areas. *Oriental Insects*, 15: 333-396.

Navajas, M., Gutierrez, J., Lagnel, J. and Boursot, P. (1996) Mitochondrial cytochrome oxidase I in tetranychid mites: a comparison between molecular phylogeny and changes of morphological and life history traits. *Bulletin of Entomological Research*, 86: 407-417.

Navajas, M., Lagnel, J., Gutierrez, J. and Boursot, P. (1998) Species-wide homogeneity of nuclear ribosomal ITS2 sequences in the spider mite *Tetranychus urticae* contrasts with extensive mitochondrial COI polymorphism. *Heredity*, 80: 742-752.

Nicastro, R.L., Sato, M.E., Arthur, V. and da Silva, M.Z. (2013) Chlorfenapyr resistance in the spider mite *Tetranychus urticae*: stability, cross-resistance and monitoring of resistance. *Phytoparasitica*, 41: 503-513.

Ochoa, R. and Aguilar, H. (1989) Combate químico de la aranita roja (*Tetranychus urticae* Koch) en fresa (*Fragaria* sp.). *Manejo Integrado de Plagas*, 11: 51-60.

Ochoa, R. and Vargas, C. (1997) Mites associated with *Epipremnum pinnatum* cv. aureum in Costa Rica. *Manejo Integrado de Plagas*, 34-43.

Ochoa, R., Aguilar, H. and Merino, F.L. (1989) Chemical control of red spider mites (Acari: Tetranychidae) in chayote (*Sechium edule* (Jacq.) SW.). *Manejo Integrado de Plagas*, 31-45.

Ohno, S., Kodama, H. and Ganaha-Kikumura, T. (2011) Temporal occurrence of the two-spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae), on Ishigaki Island, Okinawa. *Journal of the Acarological Society of Japan*, 20: 37-40.

Ohno, S., Miyagi, A., Ganaha-Kikumura, T., Gotoh, T., Kitashima, Y., Ooishi, T., Ando, T., Kijima, K., Futagami, K., Uesato, T. and Yasuda, K. (2009) Species composition of spider mites (Acari: Tetranychidae) on vegetables in Okinawa, southwestern Japan. *Applied Entomology & Zoology*, 44: 627-633.

Ohno, S., Miyagi, A., Gotoh, T., Ganaha-Kikumura, T., Shiromoto, K., Kijima, K. and Ooishi, T. (2011) Wild host plants of four spider mite species (Acari: Tetranychidae) infesting fruit crops in Okinawa. *Journal of Asia-Pacific Entomology*, 14: 281-284.

Oku, K. and Saito, Y. (2014) Do males evaluate female age for precopulatory mate guarding in the two-spotted spider mite?. *Journal of Ethology*, 32: 1-6.

Oku, K., Yano, S. and Takafuji, A. (2003) Different maternal effects on diapause induction of tetranychid mites, *Tetranychus urticae* and *T. kanzawai* (Acari: Tetranychidae). *Applied Entomology & Zoology*, 38: 225-232.

Osakabe, M., Hirose, T. and Sato, M. (2002) Discrimination of four Japanese *Tetranychus* species (Acari: Tetranychidae) using PCR-RFLP of the inter-transcribed spacer region of nuclear ribosomal DNA. *Applied Entomology & Zoology*, 37: 399-407.

Ozsisli, T. and Cobanoglu, S. (2011) Mite (Acari) fauna of some cultivated plants from Kahramanmaraş, Turkey. *African Journal of Biotechnology*, 10: 2149-2155.

Papaioannou-Souliotis, P., Markoyiannaki-Printziou, D. and Zeginis, G. (2000) Observations on acarofauna in four apple orchards of Central Greece. II. Green cover and hedges as potential sources of phytoseiid mites (Acari: Phytoseiidae). *Acarologia*, 41: 411-421.

Papaioannou-Souliotis, P., Ragusa Di Chiara, S. and Tsolakis, H. (1994) Phytophagous mites and their predators observed on cultivated plants in Greece during 1975-1990. *Annales de L'Institut Phytopathologique Benaki*, (N.S.), 17: 35-87.

Pascual-Ruiz, S., Gomez-Martinez, M.A., Ansaloni, T., Segarra-Moragues, J.G., Sabater-Munoz, B., Jacas, J.A. and Hurtado-Ruiz, M.A. (2014) Genetic structure of a phytophagous mite species affected by crop practices: the case of *Tetranychus urticae* in clementine mandarins. *Experimental and Applied Acarology*, 62: 477-498.

Perrot-Minnot, M.J., Cheval, B., Migeon, A. and Navajas, M. (2002) Contrasting effects of Wolbachia on cytoplasmic incompatibility and fecundity in the haplodiploid mite *Tetranychus urticae*. *Journal of Evolutionary Biology*, 15: 808-817.

Petrovic, A., Jurisic, A. and Rajkovic, D. (2010) Seasonal distribution and species association among spider mites (Acari: Tetranychidae) and predatory mites (Acari: Phytoseiidae and Acari: Stigmaeidae) in Serbian apple orchards. *International Journal of Acarology*, 36: 519-526.

Pozzebon, A., Duso, C., Tirello, P. and Ortiz, P.B. (2011) Toxicity of thiamethoxam to *Tetranychus urticae* Koch and *Phytoseiulus persimilis* Athias-Henriot (Acari Tetranychidae, Phytoseiidae) through different routes of exposure. *Pest Management Science*, 67: 352-359.

Rahman, K.A. and Sapra, A.N. (1940) Mites of the family Tetranychidae from Lyallpur with descriptions of four new species. *Proceedings of the Indian Academy of Science, Ser. B*, 11: 17-196.

Raworth, D.A. (2007) Initiation of oviposition after winter diapause in the spider mite *Tetranychus urticae* (Acari: Tetranychidae): prediction and historical patterns. *Population Ecology*, 49: 201-210.

Rector, B.G., Czarnoleski, M., Skoracka, A. and Lembićz, M. (2016) Change in abundance of three phytophagous mite species (Acari: Eriophyidae, Tetranychidae) on quackgrass in the presence of choke disease. *Experimental and Applied Acarology*, 70: 35-43.

Reichert, M.B., Da Silva, G.L., Rocha, M.D., Johann, L. and Ferla, N.J. (2014) Mite fauna (Acari) in soybean agroecosystem in the northwestern region of Rio Grande do Sul State, Brazil. *Systematic and Applied Acarology*, 19: 123-136.

Rigamonti, I.E. and Lozzia, G.C. (1999) Injurious and beneficial mites on urban trees in Northern Italy. *Acta Horticulturae*, 177-182.

Robles-Bermudez, A., Robles-Bermudez, G.F., Rodriguez-Maciel, J.C., C., S.-O., Lagunes-Tejeda, A., Flores-Canales, R. and Cambero Campos, J.O. (2012) Resistance of four population mites (*Tetranychus urticae* Koch.) to propargite in cut rose (*Rosa x hybrida*) in the State of Mexico, Mexico. *Revista Mexicana de Ciencias Agrícolas*, 3: 785-795.

Ros, V.I.D., Fleming, V.M., Feil, E.J. and Breeuwer, J.A.J. (2012) Diversity and recombination in Wolbachia and Cardinium from *Bryobia* spider mites. *Bmc Microbiology*, 12:.

Rossi Simons, N.H. (1961) Lista de las especies de Tetranychidae (Acari) de la Republica Argentina. Idia, 163: 9-13.

Roy, M., Brodeur, J. and Cloutier, C. (1999) Seasonal abundance of spider mites and their predators on red raspberry in Quebec, Canada. Environmental Entomology, 28: 735-747.

Saba, F. (1964) Mitteilung über Schadmilben im Vorderen Orient. Anzeiger fur Schadlingskunde Pflanzenschutz Umweltschutz, 37: 17-19.

Saba, F. (1973) Les acariens nuisibles aux plantes cultivees au Maroc. Al Awania, 49: 69-97.

Sadeghi-Namaghi, H. (2010) Mites (Acari: Prostigmata & mesostigmata) inhabiting green planting in urban environment of North-Eastern Iran, including six new records. Munis Entomology & Zoology Journal, 5: 123-130.

Sakagami, T., Saito, Y., Kongchuensin, M. and Sahara, K. (2009) Molecular phylogeny of *Stigmaeopsis*, with special reference to speciation through host plant shift. Annals of the Entomological Society of America, 102: 360-366.

Sakamoto, H. and Gotoh, T. (2017) Non-destructive direct polymerase chain reaction (direct PCR) greatly facilitates molecular identification of spider mites (Acari: Tetranychidae). Applied Entomology and Zoology, 52(4): 661-665.

Sakamoto, H., Matsuda, T., Suzuki, R., Saito, Y., Lin, J.Z., Zhang, Y.X., Sato, Y. and Gotoh, T. (2017) Molecular identification of seven species of the genus *Stigmaeopsis* (Acari: Tetranychidae) and preliminary attempts to establish their phylogenetic relationship. Systematic and Applied Acarology, 22: 91-101.

Sanchez Martinez, L., Flechtmann, C.H.W. and De Moraes, G.J. (2014) Plant mites of the Dominican Republic, with a description of a new species of *Petrobia* (Tetranychina) Waisnstein, 1960 (Acari, Prostigmata, Tetranychidae) and a key to the species of this subgenus. Zootaxa, 3846: 547-560.

Sato, M.E., Silva, M.Z.d., Raga, A. and Souza Filho, M.F.d. (2005) Abamectin resistance in *Tetranychus urticae* Koch (Acari: Tetranychidae): selection, cross-resistance and stability of resistance. Neotropical Entomology, 34: 991-998.

Sato, Y., Alba, J.M. and Sabelis, M.W. (2014) Testing for reproductive interference in the population dynamics of two congeneric species of herbivorous mites. Heredity, 113: 495-502.

Sato, Y., Alba, J.M., Egas, M. and Sabelis, M.W. (2016) The role of web sharing, species recognition and host-plant defence in interspecific competition between two herbivorous mite species. Experimental and Applied Acarology, 70: 261-274.

Sato, Y., Staudacher, H. and Sabelis, M.W. (2016) Why do males choose heterospecific females in the red spider mite?. Experimental and Applied Acarology, 68: 21-31.

Seeman, O.D. and Beard, J.J. (2011) Identification of exotic pest and Australian native and naturalised species of *Tetranychus* (Acari: Tetranychidae). Zootaxa, 2961: 1-72.

Seiedy, M., Soleymani, S. and Hakimitabar, M. (2017) Development and reproduction of the predatory mite Amblyseius swirskii Athias-Henriot (Acari: Phytoseiidae) on *Tetranychus urticae* Koch (Acari: Tetranychidae) and *Bemisia tabaci* Gennadius (Heteroptera: Aleyrodidae). International Journal of Acarology, 43: 160-164.

Seki, K. (2016) Leaf-morphology-assisted selection for resistance to two-spotted spider mite *Tetranychus urticae* Koch (Acari: Tetranychidae) in carnations (*Dianthus caryophyllus* L). Pest Management Science, 72: 1926-1933.

Sheela, K. and Ramani, N. (2012) Phytophagous mites - a potential threat to medicinal plants in Kerala, India. International Journal of Acarology, 38: 62-65.

Sheikholeslam-Zadeh, S. and Sadeghi-Nameghi, H. (2010) Injurious mites associated with broad leaf trees in Mashhad greens (In Persian). Journal of Plant Protection (Agricultural Sceince and Technology), 24: 210-217.

Shi, L., Xu, Z.F., Shen, G.M., Song, C.G., Wang, Y., Peng, J.F., Zhang, J. and He, L. (2015) Expression characteristics of two novel cytochrome P450 genes involved in fenpropathrin resistance in *Tetranychus cinnabarinus* (Boisduval). Pesticide Biochemistry and Physiology, 119: 33-41.

Sidumo, A., Langa, S., Manjate, J., Mulima, E., Dava, L. and Cossa, S. (2007) Biodiversity of spider mites (*Tetranychus* spp.) and their natural enemies in Mozambique. 8th African Crop Science Society Conference, El-Minia, Egypt, 27-31 October 2007, 1087-1089.

Sinaie, S., Namaghi, H.S. and Fekrat, L. (2018) A Multiplex PCR Assay for Simultaneous Discrimination of Two Predominant Spider Mites of the Genus *Tetranychus* (Acari: Tetranychidae) in Greenhouses of Iran. Journal of Agricultural Science and Technology, 20: 733-744.

Smiley, R.L. and Baker, E.W. (1995) A report on some tetranychid mites (Acari: Prostigmata) from Yemen. International Journal of Acarology, 21: 135-164.

Soliman, Z.R. and Mahfood, S.A. (1975) Phytophagous and predaceous mites of People's Republic of Southern Yemen. Bulletin of the Zoological Society of Egypt, 27: 78-84.

Srinivasa, N., Kavya, M.K. and Chinnamade Gowda, C. (2014) *Tetranychus turkestanii* an invasive spider mite pest of grapevine in Karnataka and Maharashtra. Ramamurthy, V.V. and Subramanian, S., Entomology as a science and IPM as a technology - the way forward, Pasighat, India: 61-62.

Stocco, R.S.M., Sato, M.E. and Santos, T.L. (2016) Stability and fitness costs associated with etoxazole resistance in *Tetranychus urticae* (Acari: Tetranychidae). Experimental and Applied Acarology, 69: 413-425.

Su, H.H., Jiang, F., Yu, M.Z., Yang, X.M., Yang, Y.Z. and Hong, X.Y. (2012) Effects of Wolbachia on rDNA-ITS2 variation and evolution in natural populations of *Tetranychus urticae* Koch. Systematic and Applied Acarology, 17: 47-52.

Suarez, A. (2004) Catalogo de acaros de la provincia de Guantanamo. Fitosenidad, 8: 23-31.

Sugasawa, J., Kitashima, Y. and Gotoh, T. (2002) Hybrid affinities between the green and the red forms of the two-spotted spider mite *Tetranychus urticae* (Acari: Tetranychidae) under laboratory and semi-natural conditions. *Applied Entomology & Zoology*, 37: 127-139.

Sun, J.-X., Guo, Y., Zhang, X., Zhu, W.-C., Chen, Y.-T. and Hong, X.-Y. (2016) Effects of host interaction with Wolbachia on cytoplasmic incompatibility in the two-spotted spider mite *Tetranychus urticae*. *Biological Journal of the Linnean Society*, n/a-n/a.

Suzuki, T., Wang, C.-H., Gotoh, T., Amano, H. and Ohyama, K. (2015) Deoxidant-induced anoxia as a physical measure for controlling spider mites (Acari: Tetranychidae). *Experimental and Applied Acarology*, 65: 293-305.

Takafuji, A. and Gotoh, T. (1999) Diapause characteristics of some *Tetranychus* (Acari: Tetranychidae) mites of Okinawa island. *Journal of the Acarological Society of Japan*, 8: 51-54.

Takafuji, A. and Hinomoto, N. (2008) The distribution and geographic variation in diapause capacity among populations of two *Tetranychus* species (Acari: Tetranychidae) in East and Southeast Asia. *Journal of the Acarological Society of Japan*, 17: 1-15.

Takafuji, A., Hinomoto, N., Shih, C.I.T., Gotoh, T., Ho, C.C. and Wang, C.C. (2005) Diapause characteristics of the Taiwanese populations of *Tetranychus kanzawai* Kishida and *T. urticae* Koch (Acari: Tetranychidae). *Plant Protection Bulletin*, 47: 103-114.

Takafuji, A., Santoso, S., Hinomoto, N., Shih, C.I.T., Ho, C.C. and Gotoh, T. (2003) Diapause characteristics of two species of tetranychid mites (Acari: Tetranychidae) in southern Japan and Taiwan. *Applied Entomology & Zoology*, 38: 267-270.

Tang, X.F., Zhang, Y.J., Wu, Q.J., Xie, W. and Wang, S.L. (2014) Stage-Specific Expression of Resistance to Different Acaricides in Four Field Populations of *Tetranychus urticae* (Acari: Tetranychidae). *Journal of Economic Entomology*, 107: 1900-1907.

Tirello, P., Pozzebon, A., Cassanelli, S., Van Leeuwen, T. and Duso, C. (2012) Resistance to acaricides in Italian strains of *Tetranychus urticae*: toxicological and enzymatic assays. *Experimental and Applied Acarology*, 57: 53-64.

Toda, S., Osakabe, M. and Komazaki, S. (2000) Interspecific diversity of mitochondrial COI sequences in Japanese *Panonychus* species (Acari: Tetranychidae). *Experimental & Applied Acarology*, 24: 821-829.

Tollerup, K.E., Marcum, D., Wilson, R. and Godfrey, L. (2013) Binomial and Enumerative Sampling of *Tetranychus urticae* (Acari: Tetranychidae) on Peppermint in California. *Journal of Economic Entomology*, 106: 1707-1715.

Tsagkarakis, A.E., Emmanouel, N.G., Panou, H.N., Kapaxidi, E.V. and Papadoulis, G.T. (2011) Composition and seasonal abundance of mites associated with Citrus in Greece. *International Journal of Acarology*, 37: 252-259.

Tseng, Y.H. (1974) Systematics and distribution of phytophagous and predatory mites on grape in Taiwan. Part I. Phytophagous mites. *Journal of the Agricultural Association of China*, 88: 56-73.

Tuan, S.J., Lin, Y.H., Peng, S.C. and Lai, W.H. (2016) Predatory efficacy of *Orius strigicollis* (Hemiptera: Anthocoridae) against *Tetranychus urticae* (Acarina: Tetranychidae) on strawberry. *Journal of Asia-Pacific Entomology*, 19: 109-114.

Tuan, S.J., Lin, Y.H., Yang, C.M., Atluhan, R., Saska, P. and Chi, H. (2016) Survival and reproductive strategies in two-spotted spider mites: demographic analysis of arrhenotokous parthenogenesis of *Tetranychus urticae* (Acari: Tetranychidae). *Journal of Economic Entomology*, 109: 502-509.

Tuttle, D.M., Baker, E.W. and Abbatiello, M. (1974) Spider mites from northwestern and north central Mexico (Acarina: Tetranychidae). *Smithsonian Contributions to Zoology*, 171: 1-18.

Tuttle, D.M., Baker, E.W. and Sales, F.M. (1977) Spider mites (Tetranychidae: Acarina) of the state of Ceara, Brazil. *International Journal of Acarology*, 3: 1-8.

Uddin, M.N., Alam, M.Z., Miah, M.R.U., Mian, M.I.H. and Kishowar, E.M. (2017) Life table parameters of an indigenous strain of *Neoseiulus californicus* McGregor (Acari: Phytoseiidae) when fed *Tetranychus urticae* Koch (Acari: Tetranychidae). *Entomological Research*, 47: 84-93.

Uesugi, R. and Osakabe, M. (2007) Isolation and characterization of microsatellite loci in the two-spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae). *Molecular Ecology Notes*, 7: 290-292.

Ullah, M.S. and Lim, U.T. (2017) Synergism of *Beauveria bassiana* and *Phytoseiulus persimilis* in control of *Tetranychus urticae* on bean plants. *Systematic and Applied Acarology*, 22: 1924-1935.

Urueta, E.J. (1975) Aranas rojas (Acarina: Tetranychidae) del Departamento de Antioquia. *Revista Colombiana de Entomología*, 1: 1-14.

Van de Bund, C.F. and Helle, W. (1960) Investigations on the *Tetranychus urticae* complex in N. W. Europe (Acari: Tetranychidae). *Entomologia Experimentalis et Applicata*, 3: 142-156.

Vangansbeke, D., Audenaert, J., Nguyen, D.T., Verhoeven, R., Gobin, B., Tirry, L. and De Clercq, P. (2015) Diurnal temperature variations affect development of a herbivorous arthropod pest and its predators. *Plos One*, 10: 19.

Vassiliou, V.A. and Kitsis, P. (2013) Acaricide Resistance in *Tetranychus urticae* (Acari: Tetranychidae) Populations From Cyprus. *Journal of Economic Entomology*, 106: 1848-1854.

Vrydaghs, J. and Cooreman, J. (1953) L'araignée rouge du coton au Congo belge. *Bull. Agric. Congo Belge*, 44: 1325-1334.

Wang, C.H., Suzuki, T., Ohyama, K., Ullah, M.S. and Gotoh, T. (2016) Anoxia treatment for selectively controlling spider mites *Tetranychus urticae* and *Panonychus citri* with little impact on the predatory mite *Neoseiulus californicus*. *International Journal of Acarology*, 42: 206-211.

Wang, J.u., Zhang, J.P., Dou, W. and Zhao, Z.M. (2008) Influence of simulated acid rain on population dynamics of carmine spider mite, *Tetranychus cinnabarinus* (Boisduval) (Acari: Tetranychidae) and its host plant. *International Journal of Acarology*, 34: 427-434.

Wang, L., Zhang, Y.J., Xie, W., Wu, Q.J. and Wang, S.L. (2016) Sublethal effects of spinetoram on the two-spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae). *Pesticide Biochemistry and Physiology*, 132: 102-107.

Wekesa, V.W., Vital, S., Silva, R.A., Ortega, E.M.M., Klingen, I. and Delalibera, I., Jr. (2011) The effect of host plants on *Tetranychus evansi*, *Tetranychus urticae* (Acari: Tetranychidae) and on their fungal pathogen *Neozygites floridana* (Entomophthorales: Neozygitaceae). *Journal of Invertebrate Pathology*, 107: 139-145.

Wilson, L.T., Pickett, C.H., Leigh, T.F. and Carey, J.R. (1987) Spider mite (Acari: Tetranychidae) infestation foci: cotton yield reduction. *Environmental Entomology*, 16: 614-617.

Witul, A. and Kielkewicz, M. (1999) Life-history parameters of two closely related forms of the *Tetranychus urtica* complex on different host plants. Bruin, J., Van der Geest, L.P.S. and Sabelis, M.W., *Ecology of the Acari*, Kluwer Academic Publishers: 399-404.

Womersley, H. (1940) Studies in Australian Acarina, Tetranychidae and Trichadenidae. *Transactions of the Royal Society of South Australia*, 64: 233-265.

Woods, J.L., Dreves, A.J., James, D.G., Lee, J.C., Walsh, D.B. and Gent, D.H. (2014) Development of Biological Control of *Tetranychus urticae* (Acari: Tetranychidae) and *Phorodon humuli* (Hemiptera: Aphididae) in Oregon Hop Yards. *Journal of Economic Entomology*, 107: 570-581.

Woolhouse, M.E.J. and Harmsen, R. (1984) The mite complex on the foliage of a pesticide-free apple orchard: population dynamics and habitat associations. *Proceedings of the Entomological Society of Ontario*, 115: 1-11.

Xie, L., Hong, X.Y. and Xue, X.F. (2006) Population genetic structure of the two spotted spider mite (Acari: Tetranychidae) from China. *Annals of the Entomological Society of America*, 99: 959-965.

Xie, L., Miao, H. and Hong, X.Y. (2006) The two spotted spider mite *Tetranychus urticae* Koch and the carmine spider mite *Tetranychus cinnabarinus* (Boisduval) in China mixed in their Wolbachia phylogenetic tree. *Zootaxa*, 1165: 33-46.

Zacharda, M., Pultar, O. and Muska, J. (1988) Washing technique for monitoring mites in apple orchards. *Experimental & Applied Acarology*, 5: 181-183.

Zaher, M.A., Gomaa, E.A. and El-Enany, M.A. (1982) Spider mites of Egypt (Acari: Tetranychidae). *International Journal of Acarology*, 8: 91-114.

Zeity, M., Srinivas, N. and Gowda, C.C. (2017) Are *Tetranychus macfarlanei* Baker and Pritchard and *Tetranychus malaysiensis* Ehara (Acari: Tetranychidae) one species? Morphological and molecular evidences for synonymy between these two spider mite species and a note on invasiveness of *T. macfarlanei* on okra and eggplant in India. *Systematic and Applied Acarology*, 22: 467.

Zhang, J., Sun, J.T., Jin, P.Y. and Hong, X.Y. (2016) Development of microsatellite markers for six *Tetranychus* species by transfer from *Tetranychus urticae* genome. *Experimental and Applied Acarology*, 70: 17-34.

Zhang, Y.C., Xu, Z.F., Wu, Q., Peng, M., Liu, Y.C., Liu, X., Shi, L., Shen, G.M., Pan, Y. and He, L. (2016) Identification of differentially expressed microRNAs between the fenpropathrin resistant and susceptible strains in *Tetranychus cinnabarinus*. Plos One, 11: 16.

Zhang, Z. and Jacobson, R.J. (2000) Using adult female morphological characters for differentiating *Tetranychus urticae* complex (Acari: Tetranychidae) from greenhouse tomato crops in UK. Systematic & Applied Acarology, 5: 69-76.

Zhao, D.-X., Chen, D.-S., Ge, C., Gotoh, T. and Hong, X.-Y. (2013) Multiple Infections with Cardinium and two strains of Wolbachia in The Spider Mite *Tetranychus phaselus* Ehara: Revealing New Forces Driving the spread of Wolbachia. Plos One, 8: e54964.

Zriki, G., Shaabo, A. and Boubou, A. (2015) A preliminary survey of the spider mites (Acari: Tetranychidae) in Latakia governorate of Syria. Acarologia, 55: 303-309.