

Table 6. Average dominance [%] of collected mite species/taxa in the particular types of farm buildings examined in Żywiecki district.

Type of farming building	<i>Acarus siro</i>	<i>Acarus farris</i>	<i>Acarus immobilis</i>	<i>Tyrophagus putrescentiae</i>	<i>Tyrophagus longior</i>	<i>Sancassania berlessei</i>	<i>Rhizoglyphus echinopus</i>	<i>Rhizoglyphus robini</i>	<i>Lepidoglyphus destructor</i>	<i>Lepidoglyphus michaeli</i>	<i>Lepidoglyphus fustifer</i>	<i>Glycyphagus domesticus</i>	<i>Glycyphagus privatus</i>	<i>Glycyphagus ornatus</i>	<i>Gohieria fusca</i>	<i>Ctenoglyphus plumiger</i>	<i>Ctenoglyphus canestrinii</i>	<i>Chortoglyphus arcuatus</i>	<i>Gymnoglyphus longior</i>	<i>Hirstia chalidonis</i>	Oribatida (sensu lato)	Cheyletidae	Cunaxidae	Heterostigmata	Prostigmata (other)	Mesostigmata	
1	47.33	25.79	2.53	16.57	0.00	0.00	0.00	0.00	15.34	3.91	0.00	10.93	0.00	0.00	2.59	9.31	6.73	8.79	1.97	3.45	0.00	21.90	0.00	0.00	0.00	0.00	6.43
2	20.09	15.99	8.05	15.47	2.44	0.00	3.00	0.00	23.49	1.45	17.71	9.72	0.00	0.72	7.74	2.67	0.00	17.85	0.00	0.00	0.00	14.94	0.00	0.00	0.00	0.00	4.69
3	24.21	20.05	5.59	5.28	1.97	0.00	0.00	0.00	14.14	5.72	4.53	19.44	0.83	0.83	0.00	1.60	0.00	2.79	1.10	0.00	0.96	8.16	0.00	0.00	0.00	0.00	12.92
4	27.95	24.66	12.93	13.05	3.95	0.00	0.00	0.00	11.46	5.01	1.31	19.48	0.00	0.00	0.00	2.88	0.00	5.01	0.85	0.00	0.00	4.18	0.00	3.60	2.54	4.00	
5	17.36	15.47	0.00	14.07	0.00	2.44	0.00	0.00	11.69	2.07	1.42	15.34	0.00	0.00	6.67	10.08	0.00	23.35	0.00	0.00	0.00	7.12	4.97	0.00	0.00	20.40	
6	34.53	26.68	0.00	22.79	0.00	0.00	0.00	1.52	8.42	3.70	0.00	3.70	0.00	0.00	4.55	4.13	0.00	5.56	0.00	0.00	0.00	9.89	0.00	0.00	0.00	4.35	

Explanations:

1 – barns

2 – pigsties

3 – cowsheds

4 – chaff-cutter buildings

5 – poultry houses

6 – pigeon houses